

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>6</sup> :

C07K 1/04, G01N 33/53

(11) International Publication Number:

WO 95/34575

A1

(43) International Publication Date: 21 December 1995 (21.12.95)

(21) International Application Number: PCT/IB95/00560

(22) International Filing Date: 13 June 1995 (13.06.95)

(30) Priority Data:

08 260,199

15 June 1994 (15.06.94)

US

(71) Applicant: S.P.I. SYNTHETIC PEPTIDES INCORPORATED  
[CA,CA]; 4465 99th Street, Edmonton, Alberta T6E 5B6  
(CA).(72) Inventors: HODGES, Robert, S.; 9045 Saskatchewan Drive,  
Edmonton, Alberta T6G 2B2 (CA). IRVIN, Randall,  
T.; 9 Chelsea Manor, Sherwood Park, Alberta T8H 1C1  
(CA). HOLM, Arne; Chemistry Dept., Royal Veterinary  
and Agricultural University, Thorvaldsensvej 40, DK-1871  
Frederiksberg (DK). WONG, Wah, Y.; #66 Bellamy Hill,  
Edmonton, Alberta T5K 0T1 (CA). SHETH, Hasmukh,  
B.; #1005, 11025-82 Avenue, Edmonton, Alberta T6G  
0T1 (CA). HUSBAND, Devon, L.; 10046-117 Street #6,  
Edmonton, Alberta T5K 1X2 (CA).(74) Agent: ERRATT, Judy, A.; Gowling, Strathy & Henderson,  
Suite 2600, 160 Elgin Street, Ottawa, Ontario K1P 1C3  
(CA).(81) Designated States: AU, CA, JP, European patent (AT, BE,  
CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT,  
SE).**Published***With international search report.**Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.*

(54) Title: COMBINATORIAL PEPTIDE LIBRARY AND METHOD

## (57) Abstract

A combinatorial library composition and method for using the library to construct oligomers effective to bind to a selected ligand is disclosed. The library composition includes at least two sets of combinatorial oligomer libraries, each library set having selected oligomer subunit positions filled by known subunits, and other subunit positions containing permutations of subunits. In the selection method, oligomers from each library set are identified, and a new permutation library formed of subunits corresponding to the highest binding affinity oligomers in each library is screened for binding affinity to the selected ligand.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## COMBINATORIAL PEPTIDE LIBRARY AND METHOD

1. Field of the Invention

The present invention relates to combinatorial peptide libraries, methods of screening  
5 such libraries and the identification of synthetic peptide antigens.

2. References

- Barbas, C.F., et al., *Proc. Natl. Acad. Sci. USA* 89(10):4457 (1992).  
Blake, J., and Litzi-Davis, L., *Bioconjugate Chem.* 3:510-513 (1992).  
10 Doig, P., et al., *Infect. Immun.* 58:124-130 (1990).  
Dooley, C.T., et al., *Proc. Natl. Acad. Sci. USA* 90(22):10822 (1993a).  
Dooley, C.T., et al., *Life Sci.* 52(18):1509 (1993b).  
Ecker, D.J., et al., *Nuc. Acids Res.* 21(8):1853 (1993).  
Eichler, J., et al., *Biochemistry* 32(41):11035 (1993).  
15 Erickson, B.W., and Merrifield, R.B., in THE PROTEINS, p. 255, Academic Press,  
Inc., New York (1976).  
Fan, N., et al., *J. Clin. Micro.* 30(4):905 (1992).  
Fields, G.B., and Noble, R.L., *Int. J. Peptide Protein Res.* 35:161-214 (1990).  
Fodor, S.P.A., et al., *Science* 251:767-773 (1991).  
20 Furka, A., et al., *14th Int'l Congress on Biochem.* 5:47 (Prague, Czechoslovakia, July  
10-15) (1988a).  
Furka, A., et al., *14th Int'l Congress on Biochem.* 5:288 (Prague, Czechoslovakia,  
August 15-19) (1988b).  
Furka, A., et al., *Int. J. Pept. Protein Res.* 37:487-493 (1991).  
25 Gausepohl, H., et al., *Peptide Res.* 5:315-320 (1992).  
Geysen, H.M., et al., *Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1984).  
Geysen, H.M., et al., *Proc. Natl. Acad. Sci. USA* 82:178-182 (1985).  
Hodges, R.S., et al., *Peptide Res.* 1:19-30 (1988).  
Holm, A., and Meldal, M., "Multiple Column Peptide Synthesis" in PEPTIDES 1988  
30 (Bayer, et al., Eds., Walter deGruyter & Co., Berlin-New York) p. 208 (1989).  
Hortin, G.L., et al., *Biochem. Int.* 26:731-738 (1992).  
Houghten, R.A., et al., *J. Biol. Chem.* 260:1551-1555 (1985).  
Houghten, R.A., et al., *Nature* 354:84-86 (1991).  
35 Houghten, R.A., et al., *BioTechniques* 13:412-421 (1992).

- King, D.S., *et al.*, *Int. J. Peptide Protein Res.* 36:255-266 (1990).
- Kramer, A., *et al.*, *Pept. Res.* 6(6):314 (1993).
- Krichbaum-Stenger, K., *et al.*, *Blood* 70:1303 (1987).
- Meldal, M., *et al.*, *Int. J. Peptide & Protein Res.* 41:250 (1993).
- 5 Ohlmayer, M.H., *et al.*, *Proc. Natl. Acad. Sci. USA* 90:23:10922 (1993).
- Pinilla, C., *et al.*, *Biotechniques* 13(6):901 (1992).
- Pinilla, C., *et al.*, *Gene* 128(1):71 (1993).
- Schulz, G.E. and R.H. Schirmer., PRINCIPLES OF PROTEIN STRUCTURE, Springer-Verlag.
- 10 Schumacher, T.N.M., *et al.*, *Eur. J. Immunol.* 22:1405-1412 (1992).
- Sebestyen, F., *et al.*, *Bioorg. Med. Chem. Lett.* 3:413-418 (1993).
- Wong, W.Y., *et al.*, *Protein Sci.* 1:1308 (1992).
- Zuckermann, R.N., *et al.*, *Int. J. Pept. Protein Res.* 40:498-507 (1992).

15 3. Background of the Invention

Currently there is widespread interest in using combinatorial libraries of random-sequence oligonucleotides, polypeptides, or synthetic oligomers to search for biologically active compounds (Kramer; Houghten, 1993a-1993c, 1992, 1991; Ohlmayer; Dooley, 1993a-1993b; Eichler; Pinilla, 1993, 1992; Ecker; and Barbas). Ligands discovered by screening libraries  
20 of this type may be useful in mimicking or blocking natural ligands, or interfering with the naturally occurring interactions of a biological target. They can also provide a starting point for developing related molecules with more desirable properties, *e.g.*, higher binding affinity.

Combinatorial libraries of the type useful in this general application may be formed by  
25 a variety of solution-phase or solid-phase methods in which mixtures of different subunits are added stepwise to growing oligomers, until a desired oligomer size is reached. A library of increasing complexity can be formed in this manner, for example, by pooling multiple choices of reagents with each additional subunit step (Houghten, 1991; 1993c).

Alternatively, the library may be formed by solid-phase synthetic methods in which  
30 beads containing different-sequence oligomers that form the library are alternately mixed and separated, with one of a selected number of subunits being added to each group of separated beads at each step. An advantage of this method is that each bead contains only one oligomer specie, allowing the beads themselves to be used for oligomer screening (Furka, 1991; Lam, 1991, 1993; Zuckermann; Sebestyen).

Still another approach that has been proposed involves the synthesis of a combinatorial library on spatially segregated arrays (Fodor). This approach is generally limited in the number of different library sequences that can be generated.

Since the chance of finding useful ligands increases with the size of the combinatorial library, it is desirable to generate libraries composed of large numbers of different-sequence oligomers. In the case of oligonucleotides, for example, a library having 4-base variability at 8 oligomer residue positions will contain as many as  $4^8$  (65,536) different sequences. In the case of a polypeptides, a library having 20-amino acid variability at six residue positions will contain as many as  $20^6$  (64,000,000) different species.

Because each different-sequence specie in a large-number library may be present in small amounts, one of the challenges in the combinatorial library selection procedure is isolating and determining the sequence of specie(s) that have the desired binding or other selected properties.

Where the combinatorial library consists of oligonucleotides, this problem may be solved by amplifying the isolated sequence, *e.g.*, by polymerase chain reaction methods. In the case of polypeptide libraries, other methods must be employed. Where the library oligomers have been formed on beads, and each bead represents one polypeptide specie, it may be possible to conduct microscale sequencing on the oligomers contained on a single isolated bead. This approach, of course, is limited to selection procedures involving particles.

In another approach, where the library has been formed by pooling multiple choices of reagents during synthesis, a pool which is shown to have desired properties is resynthesized iteratively with lower and lower complexity until a single sequence compound is identified. This approach is not only relatively laborious, but tends to "screen out" subunit permutations that may be good candidates for high-affinity oligomers, but which are eliminated at early stages of iterative resynthesis in favor of one or a few higher-affinity oligomers.

Another basic consideration in the generation of desired compounds by screening combinatorial libraries is the nature of the selected compound itself. Polynucleotide libraries are relatively easy to generate and can be sequenced at low concentrations, but the structural variation within a library is limited by the relatively few bases that are employed, typically the standard four bases/oligomer position.

In contrast to polynucleotide libraries, these also can be synthesized readily by known methods. The use of peptide libraries, however, greatly expands the potential variability of the library. As discussed above, the large number of subunit permutations in a peptide combinatorial library composed

of oligomers having 5 or more subunits sharply limits the amount of any one species available for sequence determination.

It would thus be desirable to provide a combinatorial library that retains a high potential for variability, yet can be readily manipulated to identify high-affinity binding species within the library. It is further desirable to be able to screen such a library in a solution phase, and identify likely candidates for high-binding affinity, but in a way that retains a significant number of possible subunits at each oligomer position, such that the "screening out" problem associated with reiterative resynthesis methods of sequence identification is avoided.

10 4. Summary of the Invention

The invention includes, in one aspect, a method of generating an oligomer compound capable of interacting specifically with a selected receptor.

The method includes the steps of (a) reacting the receptor with each of a set of combinatorial oligomer libraries in which one or more selected subunit positions in the library oligomers have one of the different subunit permutations in each of the selected positions, and the remaining one or more subunit positions in each library include substantially all possible combinations of the different subunits, and (b) selecting from the set of libraries, one or more libraries which show an elevated level of interaction with the receptor, thereby identifying one or more high-affinity subunit compositions corresponding to the selected subunit positions.

20 Steps (a) and (b) above are repeated with another group of selected one or more subunit positions, until substantially all of the subunit positions in the libraries oligomers have been considered. The receptor is then reacted with a permutation library of oligomers formed of placing the highest-affinity compositions identified in the steps above in each of the associated oligomer positions. From this permutation library is selected one or more oligomers which show the greatest level of interaction with the receptor.

In one example, for use in selecting a hexapeptide oligomer having three N-terminal and three C-terminal positions, one set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three N-terminal positions, and a second set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three C-terminal positions.

30 In another example, for use in selecting a hexapeptide oligomer having two N-terminal, two intermediate, and two C-terminal positions, one set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two N-terminal positions, a second set of combinatorial libraries contains a library for each of the possible permutations

of amino acids in the two intermediate positions, and a third set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two C-terminal positions.

In one general embodiment, for use in identifying a polypeptide oligomer capable of interacting specifically with a selected receptor, the subunits are representative amino acids that display the basic physico-chemical properties associated with naturally occurring amino acids, but exclude many of these naturally occurring amino acids. One group of representative amino acids include one from each of the groups (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met and Cys, (h) Pro, (i) Gln and Asn, and (j) Ser and Thr.

The method is also applicable for identifying oligomers having a large number of subunit possibilities at each oligomer position, *e.g.*, up to 50 or more subunits, such as in a library of oligomers having an upstream subunit position, and intermediate subunit position, and a downstream subunit position, one or more of which may be single-subunit positions. In such an embodiment, the combinatorial libraries used in the method include a first set of libraries containing a combinatorial library for each of the possible subunits in the upstream subunit position, a second set of combinatorial libraries containing a library for each of the possible subunits in the intermediate subunit position, and a third set of combinatorial libraries containing a library for each of the possible subunits in the downstream subunit position.

In another aspect, the invention includes a combinatorial library composition for use in selecting an oligomer compound capable of interacting specifically with a selected macromolecular receptor.

The composition includes a first set of combinatorial oligomer libraries in which one or more selected subunit positions in the library oligomers have one of substantially all possible different subunits in each of the selected positions, and the remaining one or more subunit positions in each library include substantially all possible combinations of the different subunits, and a second set of combinatorial oligomer libraries in which one or more different selected subunit positions in the library oligomers have one of substantially all possible different subunits in each of the selected positions, and the remaining one or more subunit positions in each library include substantially all possible combinations of the different subunits.

Another embodiment of the invention includes a library of oligomers having an N-terminal

the possible permutations of amino acids in the three N-terminal positions, and the set of

set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three C-terminal positions.

As another example, for use in selecting a hexapeptide oligomer having two N-terminal, two intermediate, and two C-terminal positions, the first set of combinatorial libraries  
5 contains a library for each of the possible permutations of amino acids in the two N terminal positions, and the second set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two intermediate positions. The composition further includes a third set of combinatorial libraries containing a library for each of the possible permutations of amino acids in the two C-terminal positions.

10 In one general embodiment, for use in identifying a polypeptide oligomer capable of interacting specifically with a selected receptor, the subunits are representative amino acids that display the basic physico-chemical properties associated with naturally occurring amino acids, such as described above.

In another general embodiment, the composition is designed for identifying oligomers  
15 having a large number of subunit possibilities at each oligomer position, *e.g.*, up to 50 or more subunits, such as in a library of oligomers having an upstream subunit position, and intermediate subunit position, and a downstream subunit position, one or more of which may be single-subunit positions.

In this embodiment, the composition includes a first set of libraries containing a  
20 combinatorial library for each of the possible subunits in the upstream subunit position, a second set of combinatorial libraries containing a library for each of the possible subunits in the intermediate subunit position, and a third set of combinatorial libraries containing a library for each of the possible subunits in the downstream subunit position.

The method and composition above have been used to construct novel antigens that are  
25 immunoreactive with antibodies directed against certain pathogenic bacterial and fungal organisms. The novel antigens, which form yet another aspect of the invention, are identified by SEQ ID NOS:3, 4, 5, 18, 21, 22, and 23.

These and other objects and features of the invention will be more fully appreciated when the following detailed description of the invention is read in conjunction with the accompanying drawings.

30

#### Brief Description of the Figures

Figs. 1A-1C are representations of combinatorial libraries of hexamers having different subunits  $X_i$  at positions 1-6 (Fig. 1A), and two complementary sets of libraries (Figs. 1B and 1C) formed in accordance with the invention;



Figs. 2A and 2B show representative peptide combinatorial libraries corresponding to those represented in Figs. 1B and 1C, respectively;

Figs. 3A-3D are representations of a combinatorial library of hexamers having different subunits X<sub>i</sub> at positions 1-6 (Fig. 3A), and three complementary sets of libraries (Figs. 3B-3D) formed in accordance with the invention;

Figs. 4A-4C show representative peptide combinatorial libraries corresponding to those represented in Figs. 3B-3D, respectively;

Figs. 5A-5D are representations of a combinatorial library of trimers having different subunits S<sub>i</sub> at positions 1-3 (Fig. 5A), and three complementary sets of libraries (Figs. 5B-5D) formed in accordance with the invention;

Figs. 6A and 6B illustrate the selection of four combinatorial libraries from each of two sets of complementary hexapeptide libraries such as those shown in Figs. 2A and 2B;

Fig. 7 shows representative members of a permutation peptide library formed from the amino acid sequences selected in Figs. 6A and 6B, and further selection of the permutation library;

Figs. 8A-8C illustrate the selection of eight, one, and five combinatorial libraries, respectively, from each of three sets of complementary hexapeptide libraries such as those shown in Figs. 4A-4C, respectively;

Fig. 9 shows representative members of a permutation peptide library formed from the amino acid sequences selected in Figs. 8A-8C;

Fig. 10 shows representative members of another permutation peptide library having peptides including amino acids other than the "representative" amino acids used to construct the "reduced set" libraries in Figs. 8A-8C;

Figs. 11A-11C show actual antibody binding data for the screening of the three sets of hexapeptide libraries indicated in Figs. 8A-8C, respectively;

Fig. 12 shows a portion of a microtitre plate used in screening a library set in the invention;

Figs. 13A and 13B illustrate steps in one method for screening combinatorial libraries for the presence of oligomers having specific binding affinity for an antibody receptor; and

Figs. 14A-14C illustrate steps in another method of screening libraries for the presence

Detailed Description of the InventionI. Definitions

Unless otherwise indicated, all terms used herein have the same meaning given below:

An "oligomer" refers to a polymer composed of typically between about 3-12 subunits, where each subunit may be one of a plurality of different possible subunits.

A "combinatorial library of oligomers" refers to a library of oligomer molecules containing a large number, typically between  $10^3$  and  $10^7$  different-sequence oligomers, typically defined by a different sequence of subunits. Each sequence in a library is preferably represented by a plurality, *e.g.*,  $10^8$ - $10^{12}$  molecules of the same sequence.

"Subunit position" in a combinatorial library of oligomers refers to the residue position of a subunit in a library of same-size oligomers. Where the oligomer is represented as a left-to-right linear string of subunits, the leftmost position will be designated as position 1, the position immediately to the right of position 1, as position 2, and so on.

"Permutations of different subunits" refers to known permutations of different subunits placed in selected oligomer positions. Thus, for example, several amino acid permutations of the first two positions in a oligopeptide include AlaAla, AlaArg, AlaAsn, AlaAsp, and so on.

"Combinations of different subunits" refers to random combinations of different subunits in selected oligomer positions, preferably including all or substantially all possible combinations of different subunits.

A "macromolecular receptor" or "receptor" is a macromolecule capable of specifically interacting with ligand molecules, including oligomers selected in accordance with the inventions. Binding of the ligand to the receptor is typically characterized by a high binding affinity, *i.e.*,  $K_b > 10^5 \text{ mole}^{-1}$ , and is intended to affect, *e.g.*, inhibit, the function of the receptor in its normal biological setting. The receptor is also referred to herein as a target structure.

"Representative amino acids that display the basic physico-chemical properties associated with naturally occurring amino acids" refer to a set of amino acids that include at least one amino acid for each group of amino acids whose side chain(s) confer a selected size, hydrophobicity, charge, and/or structure-forming property on the amino acid, as described in Section II below. Abbreviations for amino acid residues are the standard 3-letter and/or 1-letter codes used in the art to refer to one of the 20 common L-amino acids.

## II. Library Composition

This section describes a combinatorial library composition constructed in accordance with the invention.

The composition includes two or more sets of combinatorial oligomer libraries. In each set, one or more selected subunit positions have one of substantially all possible different subunits in each of the selected positions, and the remaining one or more subunit positions include substantially all possible combinations of the different subunits.

Figs 1A-1C illustrate a general a combinatorial library composition, where the oligomer is a hexamer, and the composition includes two library sets. The hexamer in Fig. 1A contains six subunits "X" at subunit positions 1-6. The subunits forming the oligomer may be, for example, amino acids in an oligopeptide, including standard and non-standard and derivatized L- and D-amino acids, amino acids coupled through non-amide linkages, forming an polypeptoid, nucleotides or trinucleotide codons in a polynucleotide, or other subunits based on ribose, glucose, or modified sugars with appropriate side chains or bases. Alternatively, the oligomer may be a molecular structure having a plurality of "subunit" R groups attached to the structure at designated positions on the structure, *e.g.*, six different substitutions position on a parent cyclic structure.

For purposes of illustration, the library composition will be described with respect to a hexapeptide composition in which the allowed subunits include some or all of the standard L-amino acids. The first library set, illustrated in Fig. 1B, is formed by filling each of the first three positions, indicated by  $O_1$ ,  $O_2$ , and  $O_3$ , with each of the possible subunits. Each of the  $O_1O_2O_3$  permutations will then form one library in the set, with the remaining positions  $X_4$ ,  $X_5$ , and  $X_6$  in each library preferably containing all or substantially all combinations of the different subunits placed in the subunits positions 4-6.

Representative members of this library set are shown in Fig. 2A, where each library contains one of the 8,000 possible three-subunit permutations in its first three position, such as permutations AAA, AAR, ARA, and so on to VVV. The last three positions, *i.e.*, positions 4-6 in each of these 8,000 libraries, preferably contain all or substantially all of the 8,000 combinations of the different subunits. That is each library in a set includes a known sequence of subunits at positions 1-3, and a combinatorial library of sequences at positions 4-6.

Each possible subunit combination for the remaining positions is included in each library preferably

containing all or substantially all combinations of the different subunits placed in subunit positions 1-3.

Representative members of this library set are shown in Fig. 2B, where each library contains one of the 8,000 possible three-subunit permutations in its last three position, such as permutations AAA, AAR, ARA, and so on to VVV. The first three positions, *i.e.*, positions 1-3 in each of these 8,000 libraries preferably contain all or substantially all of the 8,000 combinations of the different subunits. That is, each library in a set includes a known sequence of subunits at positions 4-6, and a combinatorial library of sequences at positions 1-3.

The two library sets are complementary in that together, they include a combinatorial library for each of a known subunit permutation at positions 1-3 and 4-6.

In one preferred embodiment, which is useful for purposes of simplifying the synthesis and selection procedures in the method of the invention, the subunits forming the oligomers in the library composition are representative amino acids, typically including between about 8-12 amino acids that are representative of most of all of the commonly classified groups of amino acids, based on the physico-chemical properties of the amino acids. These properties include the size, hydrophobicity, charge, and/or structure-forming properties that the side chains impart on the amino acid. Libraries generated using such a reduced set of representative amino acids are termed reduced combinatorial peptide libraries, or RCPLs.

One recognized grouping of amino acids by physico-chemical properties includes the groups (a) Ala, representing a small, uncharged side chain; (b) Glu and Asp, representing negatively charged amino acids; (c) Phe, Tyr, and Trp, representing side chains with aromatic groups; (d) Gly, representing a very small side chain and one which confers high flexibility of backbone conformation, (e) Ile and Val, representing  $\beta$ -branched, hydrophobic side chains; (f) Lys, His, and Arg, representing positively charged amino acids; (g) Leu, Met, and Cys, representing large hydrophobic and sulfur-containing side chains; (h) Pro, representing a side chain with a strong influence on secondary structure, and in particular, on turns; (i) Gln and Asn, representing amide-containing side chains; and (j) Ser and Thr, representing hydroxyl-containing side chains.

One exemplary set of amino acids include the ten amino acids Ala (A) from group (a); Glu (E) from group (b); Phe (F) from group (c); Gly (G) from group (d); Ile (I) from group (e); Lys (K) from group (f); Leu (L) from group (g); Pro (P) from group (h); Gln (Q) from group (i); and Ser (S) from group (j).

As can be appreciated, this group of amino acids reduces the number of libraries in the three-position library sets described with reference to Figs. 1 and 2 from 8,000 to 1,000 (10

$\times 10 \times 10$ ), and similarly reduces the number of combinations in each combinatorial library from 8,000 to 1,000. That is, each of the 1,000 libraries in each set includes now only 1,000 different sequences in positions 4-6, for the first library set, and at position 1-3, for the second library set.

5 Figs. 3A-3D illustrate another general combinatorial library composition, where the oligomer is a hexamer, and the composition includes three library sets. The hexamer in Fig. 3A contains six subunits "X" at subunit positions 1-6. The subunits forming the oligomer may be, for example, any of the compositions described above in reference to Figs. 1A-1C. One embodiment, described below, is comprised of three sets of hexapeptide libraries.

10 The first library set, illustrated in Fig. 3B, is formed by filling each of the first two positions, indicated by  $O_1$  and  $O_2$ , with each of the possible subunits. Each of the  $O_1O_2$  permutations will then form one library in the set, with the remaining positions  $X_3$ ,  $X_4$ ,  $X_5$ , and  $X_6$  in each library preferably containing all or substantially all combinations of the different subunits placed in the subunits positions 3-6.

15 Representative members of this library set are shown in Fig. 4A, where each library contains one of the 400 possible two-subunit permutations in its first two position, such as permutations AA, AR, RA, and so on to VV. The last four positions, *i.e.*, positions 3-6 in each of these 400 libraries preferably contain all or substantially all of the 160,000 combinations of the different subunits at positions 3-6. That is, each library in a set includes  
20 a known sequence of subunits at positions 1 and 2, and a combinatorial library of sequences at positions 3-6.

The second set of libraries in the composition is shown at Fig. 3C. Each library in this set is formed by filling the intermediate two positions, indicated by  $O_3$  and  $O_4$ , with each of the possible subunits, with the remaining positions  $X_1$ ,  $X_2$ ,  $X_5$ , and  $X_6$  in each library  
25 preferably containing all or substantially all combinations of the different subunits placed in subunit positions 3 and 4.

Representative members of this library set are shown in Fig. 4B. Each library contains one of the 400 possible two-subunit permutations in its intermediate two positions, and preferably all or substantially all of the 160,000 combinations of different subunits at positions  
30 1, 2, 5, and 6.

The third set of libraries in the composition is shown at Fig. 3D. Each library in this set

substantially all of the 160,000 possible combinations of subunits at positions 1-4 in each of the libraries in the set, as illustrated in Fig. 4C.

The three library sets are complementary in that together, they include a combinatorial library for each of a known subunit permutation at positions (1,2), (3,4), and (5,6).

Another type of combinatorial library composition includes oligomers composed of subunits in which the subunits may be, for example, backbone moieties, such as peptide or ribose, deoxyribose, or glucose, with a multiplicity (*e.g.*, 20-100) R group side chains. Alternatively, the subunits may be, for example, a single parent molecule with three "R" positions, substituted at any of the three R positions. The R groups here may be any of a large number of common organic-compound radicals, such as alkyl groups, substituted alkyl groups, aryl groups, substituted aryl groups, and acid, aldehyde, amine, sulfhydryl, and alcohol groups, either alone or in combination with alkyl or aryl groups.

An example of this type of composition is illustrated in Figs. 5A-5D. In this case, the oligomer is a trimer (Fig. 5A), which contains three subunits "S" at subunit positions 1-3, where  $S_1$  has one of a plurality of possible R groups, indicated by  $R_i$ , subunit  $S_2$  carries one of a plurality of R groups indicated by  $R_j$ , and so forth.

The first library set, illustrated in Fig. 5B, is formed by placing at the  $R_i$  position, each of  $n$  different R groups. Each subunit  $S_1$  having a different R group will then form one library in the set, with the remaining subunits in each library preferably containing all or substantially all combinations of the different R groups for  $R_j$   $R_k$ . That is, the composition includes  $N$  libraries which have a known R group in the first position and are combinatorial in the second and third positions.

Similarly, a second set of  $n$  combinatorial libraries will have  $n$  different known R group at the  $R_j$  positions and be combinatorial in  $R_i$ ,  $R_k$ , as shown in Fig. 5C, and similarly for a third set, shown in Fig. 5D.

The three library sets are complementary in that together, they include a combinatorial library for each of a known subunit R-group permutation at positions 1, 2 and 3.

### III. Construction of Libraries

This section describes the synthesis of a composition of library sets, as described above, in which the subunits are peptides formed by a representative group of L-amino acids.

As noted above, both solution-phase and solid-phase synthetic methods may be employed. In one general embodiment for producing one of the libraries in a library set, the solid-phase resin used in the method is reacted with either a mixture of subunits, at the "combinatorial" subunit positions, or with a known monomer or dimer subunit, at the "known" subunit positions.

By way of example, to prepare a library set of 100 combinatorial hexapeptide libraries (using 10 representative amino acids only), in which the first two positions are known, and the remaining four positions are combinatorial, one would distribute resin into 100 different reaction wells, and form the 100 different permutations of amino acids at the first two subunits  
5 formed on the resin. Thereafter, each reaction well would be reacted stepwise with a mixture of the 10 different subunits, effectively placing each of the possible subunits at the third position in each of the 100 libraries. This latter reaction involving a mixture of all amino acids is then repeated for the fourth, fifth, and sixth subunits positions, to form the desired library set. The general reaction chemistry may follow standard methods (*e.g.*, Holm, Meldal)

10 A second library set having known subunit positions in the third and fourth position is similarly formed by first carrying out the first and second coupling reactions with amino acid mixtures in a single vessel, carrying out the third and fourth coupling reactions in 100 separate wells with different known combinations of two subunits in each well, and the fifth and sixth reactions with the amino acid mixtures again, but in the separate wells. The final library set  
15 with known permutations of amino acids at the fifth and sixth positions is similarly formed, carrying out the first four position additions in a single vessel with amino acid mixtures, and the final two additions in separate wells.

In another general approach, in which combinatorial libraries are formed in discrete substrates, such as pins, the different pins could contain each library of a set. Thus, each of  
20 the three library sets just described could be formed on a set of 100 pins, again employing known solid-phase coupling methods (Geysen, 1984; Geysen, 1985).

The general approach outlined above was followed to construct three sets of combinatorial libraries for hexapeptides constructed of ten representative amino acids, as detailed in Example 1. Synthesis of the library sets was on NovaSyn PR 500 resin  
25 (Novabiochem, La Jolla, CA), using Fmoc amino acid Pfp-esters (Novabiochem) and OtBu (Glu), tBu (Ser) and tBoc (Lys) side-chain protecting groups, as detailed in Example 1. Other resins amenable for use on an MCPSI include Rink-AM resin (Aminotech, Nepean, ON, Canada) and Benzhydrylamine resin (Bachem, Torrance, CA).

At the known (selected) subunit positions in each peptide mixture, a 10 fold excess of  
30 the Fmoc amino acid Pfp-esters in DMF was used. At random positions, a mixture of 0.1

combinatorial libraries for use in the application of the invention to the formation of hexapeptides.

subunits, following conventional solution-phase or solid-phase synthetic methods for those different subunits.

Similarly, if the library sets are formed by substitution of different R groups on a parent structure, as described with reference to Figs. 5A-5D, the synthetic scheme for selectively placing R groups at each of the different substitution positions may employ orthogonal protection groups, allowing deprotection and substitution at each position successively. Suitable orthogonal protection groups, and methods for deprotecting them for R-group coupling, are well known.

#### 10 IV. Method of Selecting High-Affinity Oligomers

This section describes the method of the invention for generating an oligomer compound capable of interacting specifically with a selected macromolecular receptor, employing the combinatorial library composition described in Section II above.

In the method, each library in the library set is screened for its ability to interact specifically with a selected receptor, typically a macromolecular receptor. This reaction is typically a binding reaction, as measured by the formation of a binding complex between the receptor and one or more molecules in the library being screened.

The receptor is any biological receptor of interest, that is, one for which it is desired to identify an oligomer (ligand) that binds specifically to the receptor, to affect the functioning of the receptor in its normal physiological setting.

For example, the receptor may be an enzyme, where the oligomer is able to bind to the active site of the enzyme or otherwise inhibit or affect the action of the enzyme on a normal substrate.

In another general embodiment, the receptor may be a cell receptor protein, such as an ion channel or other transport receptor protein, or a receptor site for a hormone or other cell effector, or a receptor site for binding of pathogenic bacteria or viruses to a cell surface. The receptor protein may be associated with isolated cells with culture cells, with biological membrane particles isolated from tissues, with cells which are transformed to produce the receptor recombinantly, or with isolated cell receptors. Receptor proteins of this type, and expressed or isolated in a variety of forms, have been described in the literature, such as that cited above.

In a related embodiment, the receptor is an antibody or antibody fragment, where it is desired to identify an "artificial" epitope ligand that binds specifically and with high affinity to the antibody.



In a typical application, the library of oligomers is screened for oligomer (ligand) molecules that bind specifically and with high affinity, *e.g.*, with a binding constant  $K_B$  greater than  $10^5 M^{-1}$ , to the receptor. In one embodiment, illustrated in Fig. 12, each library is screened in one of the wells in a microtitre plate, such as wells 20, 22 in plate 24. Figs. 13A and 13B show individual wells 10, 22 enlarged cross-sectional view. In the screening method, one of more resin particles of beads, such as bead 26, are placed in the wells, with each well containing one of the combinatorial libraries in a library set. That is, each bead in a selected well is coated with oligomers, such as those indicated at 29 on bead 26, representing an entire combinatorial library having fixed, known subunits in selected positions, *e.g.*, the first two oligomer positions. Thus, in a 96-well microtitre plate, 96 libraries in a set of combinatorial can be screened.

After distributing each library in a set to one of the wells in a plate, the receptor, such as antibody receptor 30 in the figures is added each well, and the reaction mixtures are incubated under conditions that allow binding to library oligomers that show high-affinity with the receptor. In Fig. 13A, the library in well 20 carried on bead 26 does not contain such binding oligomers, and the antibody remains in solution. The library carried on bead 28 in well 22 does, however, have oligomers, indicated at 32, 34, which bind the antibody receptor, as indicated.

After this incubation step, the wells are washed to remove unbound material, and the beads in each well are then incubated with a reporter-labeled antibody, such as antibody 36, which is immunoreactive with the receptor antibody. After removing the second-antibody reagent, the wells are then examined for the presence of reporter label, indicating that the bead(s) in a labeled well contain at least one oligomer that has high binding affinity for the receptor. This is seen in Fig. 12, where beads in well 22 are shown as being reported labeled.

Figs. 14A-14C illustrate another method for screening combinatorial libraries, in practicing the method of the invention. In this method, the combinatorial libraries in each set of libraries are screened in solution phase. The solution-phase libraries may be prepared, as above, with oligomer cleavage from a solid-phase resin following synthesis. Fig. 14A shows two wells 40, 42 in a microtitre plate used in the testing method. Each well in the plate, such as well 40, includes one of the combinatorial libraries to be tested, where each library is made

Each library is preincubated with a low molar concentration of an receptor, in this example, an antibody, as indicated at 50. The antibodies are immunoreactive with one or more oligomer species, such as indicated at 52, in the library in well 40, but not with the library in well 42.

- 5       The reaction mixture from each library is then added to a new well on a second plate, one library per well. As seen in Fig. 14B, the second wells, such as wells 54, 56 in plate 58, are each coated with antigen molecules, such as indicated at 60. These antigens are also immunoreactive with the receptor antibody. The added libraries are now allowed to react with the surface-bound antigen under conditions allowing immune complex formation between  
10   receptor antibody molecules are surface-bound antigen. As can be appreciated from Figs. 14B, formation of an antibody-oligomer immunocomplex prevents antibody binding to the well surface (well 54). Conversely, in the absence of such complex formation, antibody becomes bound at maximum levels to the well surface (well 56).

- Following this binding step, the wells are washed to remove unbound material, and  
15   reacted with reporter-labeled antibody, such as indicated at 62 in Fig. 14C. In wells containing bound receptor antibody, such as well 56 in Fig. 14C, the reporter-labeled antibody becomes bound to the well surface through the receptor antibody, as indicated. After washing the wells to remove unbound antibody, the wells are analyzed for the presence of bound reporter. The library or libraries in a set which are selected by this method are those that show  
20   low levels or the absence of bound reporter. In the method of the invention, the screening steps just described are repeated for each of the library sets, *e.g.*, the two sets of 1000-library sets described with respect to Figs. 2A and 2B. For each set, one or more libraries showing high affinity binding to the receptor are identified. For example, in Fig. 6A, a first library set having known amino acid subunits in the first three positions is screened, and libraries having  
25   upstream-position sequences GAY, GVW, AAW, and VLY are identified as having high-affinity antigens. These results indicate that some of the combinatorial peptides having these upstream sequences are immunoreactive with the antibody receptor.

- Similarly, the second library set is screened, and libraries having downstream-position sequences SHG, THA, TKG, and TKA are identified as having high-affinity antigens. These  
30   results indicate that some of the combinatorial peptides having these downstream sequences are immunoreactive with the antibody receptor.

Having identified in this manner, high-affinity subunits for all of the positions in the library oligomers, the next step in the method is to construct a permutation library of the high-affinity subunits. As an example, the four upstream and four downstream sequences identified

in Figs. 6A and 6B, respectively, can be combined to form 16 permutations, where the relative positions of the identified subunits are preserved.

That is, "high-affinity" upstream subunits are placed in upstream positions in the permutation oligomers, and "high-affinity" downstream subunits are placed in corresponding downstream positions. The resulting permutation library for the Fig. 6 sequence is shown in Fig. 7.

Each member of the permutation library is then tested to identify the highest affinity oligomers in the library. This may be done, for example, by determining the binding affinity of each library oligomer for the receptor, according to standard methods.

10 The method described above is illustrated in detail in Example 3, which describes a competitive ELISA screen of three complimentary sets of combinatorial libraries synthesized as detailed in Example 1. The libraries are of the form illustrated in Figs. 3A-3D; that is, three sets of hexamer libraries with two "known" adjacent subunits (a dimer) in each set. The N-terminal set (HS1) has known residues at positions 1 and 2, as indicated in Fig. 3B by O<sub>1</sub>O<sub>2</sub>,  
15 the intermediate set (HS2) at positions 3 and 4 (O<sub>3</sub>O<sub>4</sub>), and the C-terminal at positions 5 and 6 (O<sub>5</sub>O<sub>6</sub>).

Combinatorial library sets HS1, HS2 and HS3 were initially screened for their ability to inhibit the interaction of a receptor, monoclonal antibody (Mab) PK99H, with its antigen (PAK peptide, SEQ ID NO:1). The antibody was generated as described in Example 2. It  
20 binds specifically to the C-terminal receptor binding domain of the PAK pilin protein (Doig, *et al.*, 1990). The epitope recognized by PK99H consists of the hexapeptide sequence represented by SEQ ID NO:2 (Wong, 1992).

The antigen (PAK peptide, SEQ ID NO:1) was coated on the wells of an assay dish (96-well plate). The libraries in sets HS1, HS2 and HS3 were preincubated with PK99H to  
25 produce an immunocomplex between the antibody and any high-affinity library oligomers. The incubate was then added to a well on the plate, and further incubated under conditions sufficient to allow antibody binding to the surface-bound antigen.

Example of the data from such a screen are illustrated in Figs. 11A-11C. Fig. 11A shows the percent inhibition of PK99H binding to PAK (horizontal bars) obtained with HS1  
30 libraries having the amino acids indicated along the y-axis at positions 1 and 2. The vertical, dashed line indicates the 50% inhibition level. The horizontal line at the top indicates 100% inhibition.

percent inhibition of PK99H binding to PAK (horizontal bars) obtained with HS1 libraries having the amino acids indicated along the y-axis at positions 1 and 2. The vertical, dashed line indicates the 50% inhibition level. The horizontal line at the top indicates 100% inhibition.

Nine positive HS1 libraries were identified. The sequence at the first two positions (first dimer), in order of most inhibitory to least inhibitory, is EQ, QQ, QA, AF, GQ, SL, EF, GL and QL.

A similar screen is performed with sets HS2 and HS3. The results are shown in Figs. 11B and 11C, respectively. Only one positive HS2 library was identified, having the amino acids FI at positions 3 and 4 (second dimer), respectively. Five positive HS3 libraries are identified, having the sequences PK, IG, PL, LG and IL (in order of most inhibitory to least inhibitory) at positions 5 and 6 (third dimer).

A set of 15 permutation peptides was constructed, as above, using the five best dimer sequences identified in the screen of HS1 at the first dimer position, FI at the second dimer position and the three best HS3 dimer sequences at the third dimer position. The peptides were synthesized using the 10 representative amino acids indicated above, following the method detailed in Example 4.

Six of these peptides (indicated in Table 1) were soluble in aqueous medium, and were screened using a competitive ELISA, as detailed in Example 5. This assay is similar to that described above, and in Example 3, except that 10 mM, rather than 50 mM PBS is used during the incubations and the concentration of the hexapeptide being assayed is varied to construct a competitive binding curve from which  $I_{50}$  values (the concentration of peptide required to produce a 50% inhibition of antibody binding to immobilized antigen) can be determined.

The results of the screen are shown in Table 1 as  $I_{50}$  values for the six water-soluble peptides. The  $I_{50}$  for the two most effective peptides is 1.3  $\mu$ M. One of these, EQFIPK (SEQ ID NO:3) has a sequence that is contained in the sequence of the native epitope (SEQ ID NO:2). Interestingly, the sequence of EQFIPK corresponds to the dimer sequences identified as the most inhibitory in the screens of HS1, HS2 and HS3.

Two other permutation peptides, QQFIPK (SEQ ID NO:4) and QAFIPK (SEQ ID NO:5), have  $I_{50}$  values similar to that of the native sequence (1.3  $\mu$ M and 2.0  $\mu$ M respectively). Peptides GQFIPK (SEQ ID NO:7) and EQFIIG (SEQ ID NO:8) show an 8-fold and 35-fold increase in  $I_{50}$  values, respectively. One peptide, AFFIPK (SEQ ID NO:6), shows a large loss of inhibitory activity ( $I_{50}$  = 1100  $\mu$ M). The three peptides having  $I_{50}$  values of 2  $\mu$ M or less are identified by asterisk symbol in Fig. 9, and correspond to SEQ ID NOS: 3-5.

Table 1

Peptide	SEQ ID NO.:	I <sub>50</sub> (μM)
EQFIPK	3	1.3
QQFIPK	4	1.3
QAFIPK	5	2.0
AFFIPK	6	1100
GQFIPK	7	10
EQFIIG	8	45

To identify additional high-affinity binding compositions, the four highest-affinity permutation peptides identified above (SEQ ID NOS:3, 4, 5 and 7) are used as templates for the synthesis of variant permutation peptides having same-type amino acid substitutions at selected subunit positions. Twelve variants, identified by SEQ ID NOS: 18-29, were synthesized in a manner similar to the permutation peptides (detailed in Example 4), except that naturally-occurring amino acids other than the representative amino acids were used at certain selected subunit positions.

The strategy followed in the synthesis of the 12 variant permutation peptides is apparent from an examination of Table 2, which shows the sequences of the template peptides (SEQ ID NOS: 3, 4, 5, and 7) along with their respective variants peptides (SEQ ID NOS 18-29). The subunit positions substituted with same-type amino acids (selected from the 20 naturally occurring amino acids) are underlined in the sequences shown in Table 2.

The variant peptides were screened with a competitive ELISA as described above for the permutation peptides, and I<sub>50</sub> values are determined (data in Table 2). Two of the variant peptides (SEQ ID NOS:21 and 22) exhibit a higher degree of MAb PK99H antibody binding inhibition than the native sequence (0.9 μM and 1.1 μM, respectively).

Table 2

Peptide	SEQ ID NO:	ID <sub>50</sub> (μM)
EQFIPK	3	1.3
<u>D</u> QFIPK	18	2.3
EN <u>F</u> IPK	19	190
EQ <u>Y</u> IPK	20	190
EQ <u>W</u> IPK	21	0.9
EQF <u>M</u> PK	22	1.1
EQF <u>V</u> PK	23	3.4
EQFIP <u>H</u>	24	620
EQFIP <u>R</u>	25	>730
QQFIPK	4	1.3
<u>N</u> QFIPK	26	12
<u>Q</u> NFIPK	27	990
QAFIPK	5	2.0
<u>N</u> AFIPK	28	14
GQFIPK	7	10
<u>G</u> NFIPK	29	>840

It will be appreciated from the foregoing that other methods may be employed for selecting, from a set of combinatorial libraries, those libraries which show an elevated level of interaction with the ligand, in order to identify high-affinity subunit compositions.

One example is a screen based on the ability of library compositions of the present invention to inhibit the binding of a pathogen, such as a virus or bacterium, host cells. The host cells are isolated, for example, by "FICOLL-HYPAQUE" sedimentation, pelleted, resuspended in medium and incubated with a mixture of, for example, reporter-labelled virus (*i.e.* rhodamine-18-labeled virus; Fan, *et al.*, 1992) that had been pre-incubated with a combinatorial library.

The cells are then washed, fixed, for example, with 1% paraformaldehyde, and the amount of bound virions determined by, for example, flow cytometric analysis. Libraries effective to reduce the binding of virus to the cells are identified as

containing oligomers which bind with high affinity to a ligand (*i.e.* cell surface molecule) on the virus, and in so doing, block the binding of the virus to the cells.

From the foregoing, it will be appreciated how various objects and features of the invention are met. In practicing the method, the user is provided with typically two or three sets of combinatorial libraries. After screening the individual members of the library sets, one or more possible optimal subunits are identified for each subunit position. These identified subunits are then used to construct a relatively small number of permutations oligomers that can then be tested for optimal activity.

The method thus avoids the need to iteratively back synthesize new libraries at each successive screening stage, and therefore both faster and less expensive in terms of library constructs. The method also avoids "locking out" promising subunit possibilities occurs with iterative back synthesis, and thus more permutations of optimal combinations of subunits are preserved. This advantage can be appreciated from the peptides selected in Table 1 and 2 which show several high-affinity peptides produced by the method of the invention having alternative amino acid residues in each of the first four oligomer positions.

The following examples illustrate, but in no way are intended to limit the present invention.

#### MATERIALS AND METHODS

Abbreviations used: BSA, bovine serum albumin; DCM, dichloromethane; DMF, *N,N*-dimethylformamide; ELISA, enzyme-linked immunosorbent assay; Fmoc, 9-fluorenylmethoxycarbonyl; HPLC, high performance liquid chromatography;  $I_{50}$ , concentration of peptide required to produce 50% inhibition in competitive ELISA; MCPSI, Multiple Column Peptide Synthesis instrument; PAK, *Pseudomonas aeruginosa* strain K; PBS, phosphate-buffered saline; Pfp, pentafluorophenyl; RCPL, representative combinatorial peptide libraries; SPPS, solid-phase peptide synthesis; tBoc, *tert*-butoxycarbonyl; tBu, *tert*-butyl; TFA, trifluoroacetic acid.

#### EXAMPLE 1

Novabiochem, La Jolla, CA) for continuous flow synthesis of peptide amides, using a Multiple Column Peptide Synthesis Instrument (MCPSI). The physical layout and

mechanical operation of the MCPSI has been previously described in detail by Holm *et al.* (Holm, *et al.*, 1989; Meldal, *et al.*, 1993). Briefly, the instrument is a manually operated apparatus for parallel solid-phase peptide synthesis. It is constructed with a Teflon reaction block with a 96-well ( $8 \times 12$ ) layout or a 100 well ( $10 \times 10$ ) layout. Each well resembles a column in the continuous flow version of the polyamide peptide synthesis method. Two washers are installed for dispensing of solvents and deprotecting reagents in a parallel fashion. Activated and protected amino acids are transferred from a dispenser tray as solutions, eight at a time. Nitrogen gas was used to maintain a positive pressure in the chamber.

To each well, 20 mg of the resin was added. Fmoc amino acid Pfp-esters (Novabiochem, La Jolla, CA) were used for the synthesis. The side-chain protecting groups used were: OtBu (Glu), tBu (Ser) and tBoc (Lys). For those defined positions ( $O_nO_{n+1}$ ) in each peptide mixture, 0.3 M (10 folds excess) of the Fmoc amino acid Pfp-esters in DMF (200  $\mu$ l/well) were used. Double couplings were performed for 2 hr and 30 min, respectively. In the case of  $X_n$  position, a mixture of 0.1 equivalent of each of the 10 Fmoc amino acid Pfp-esters in DMF were used and coupled for 2 hr, followed by a second coupling for an additional 2 hr.

Fmoc deprotection was carried out with 20% piperidine in DMF, and acetylation of the  $\alpha$ -amino group was performed with excess acetic anhydride. The peptides were cleaved from the resin with 95% TFA/water ( $4 \times 300 \mu$ l). The TFA solution was then evaporated to dryness using a Savant "SPEEDVAC" (Savant Instruments Inc., Farmingdale, NY). The semi-solid products were washed three times with diethyl ether and then dried cautiously with a slow stream of nitrogen. The peptides were dissolved in water and centrifuged, the supernatant was transferred into a cryo-tube (Nunc, Denmark).

The synthesis was repeated until all libraries were synthesized. The peptide libraries were stored in lyophilized form.

## EXAMPLE 2

### PRODUCTION OF MONOCLONAL ANTIBODY PK99H

The monoclonal antibody, PK99H, was prepared as described by Doig, *et al.* (Doig, *et al.*, 1990). Ascites tumors were produced by injecting  $10^6$  hybridoma cells into pristine primed BALB/c male mice. Ascites fluid was collected and partial purification was accomplished with ammonium sulfate fractionation and subsequently



dialyzed against phosphate buffered saline (PBS), pH 7.4. Antibody was then purified by HPLC on a protein G affinity column (Chromato-Chem, Missoula, Montana). PK99H is an IgG subclass 1 with k light chains.

5

### EXAMPLE 3

#### COMPETITIVE ELISA SCREEN OF PEPTIDE LIBRARY

The peptide libraries were screened by competitive ELISA (Enzyme Linked Immunosorbent Assay) in 96-well microtiter plates purchased from Costar Corp. (Cambridge, MA). Microtiter plates from several other suppliers were also evaluated, but did not work as well with the peptide-antigen system described herein.

A microtiter plate was coated with the intrachain disulfide-bridged (oxidized) *Pseudomonas aeruginosa* strain K (PAK) 17-residue peptide (SEQ ID NO:1, 200  $\mu$ M, 80  $\mu$ l/well) in 0.01 M carbonate buffer, pH 9.5. This peptide antigen corresponds to the C-terminal region of the PAK pilin.

The plate was incubated for 6 hr at room temperature and was subsequently blocked with 3% bovine serum albumin (BSA) in PBS at 4°C overnight. Peptide library solutions (final concentration: 8 mM) were preincubated for an hour with the monoclonal antibody PK99H (250  $\mu$ g protein/ml in stock solution, final dilution in each well was  $2 \times 10^{-5}$  of the stock solution) in 50 mM PBS containing 0.05% BSA, pH 7.4 (incubation buffer). The solution mixtures (80  $\mu$ l) were added into corresponding wells and incubated for two hours at 37°C. The plate was washed five times with the incubation buffer.

The amount of antibody bound was detected by goat anti-mouse IgG conjugated to horseradish peroxidase (Jackson Laboratories, CA) using 2,2'-azino-di-(3-ethylbenzthiazoline sulfonic acid) as a substrate. The substrate solution was prepared in 10 mM sodium citrate buffer, pH 4.2, containing 0.03% (v/v) hydrogen peroxide. The reaction was stopped by 4 mM sodium azide, and the absorbance was measured at 405 nm by a Titertek Multiskan Plus MK II microplate reader (Flow Lab Inc., McLean, Virginia).

30

### EXAMPLE 4

Peptide libraries were synthesized following the general procedure for solid phase synthesis as described by Erickson and Merrifield (Erickson and Merrifield, 1976)

with modifications made by Hodges, *et al.* (Hodges, *et al.*, 1988) on an Applied Biosystems 430A solid phase peptide synthesizer (Foster City, CA). All amino acids used were protected at the  $\alpha$ -amino position with the *tert*-butoxycarbonyl (tBoc-) group (Bachem Inc., Philadelphia, PA). Syntheses of peptide amides were carried out using co-poly (styrene, 1% divinylbenzene) benzhydrylamine-hydrochloride resin at a substitution of 0.92 mmol amino groups/g of resin (Bachem Inc., Philadelphia, PA). All amino acids were double coupled using dicyclohexylcarbodiimide generated symmetric anhydrides in DMF for the first coupling and dichloromethane (DCM) in the second coupling. Final acetylation was performed on the instrument using acetic anhydride.

After synthesis, peptides were cleaved from the resin support with anhydrous hydrogen fluoride (20 ml/g of peptide resin) in the presence of 10% (v/v) anisole and 1% (v/v) 1,2-ethanedithiol as scavenging reagents for 1 hour at  $-4^{\circ}\text{C}$  using type 1B HF-Reaction Apparatus (Peninsula Laboratories Inc., Belmont, CA). The solvent mixture was then removed under reduced pressure. The resin was washed with anhydrous diethyl ether ( $3 \times 25$  ml), and peptide was extracted with neat acetic acid ( $3 \times 25$  ml). The peptide solution was diluted with distilled water and then lyophilized.

Purification of crude peptides was performed by reversed-phase HPLC on an Applied Biosystems 400 solvent delivery system and a 783A programmable absorbance detector connected to a Synchropak RP-4 ( $250 \times 21.2$  mm I.D.) reversed-phase column (Synchrom Inc., Lafayette, IN) using a linear AB gradient of 0.2% B/min at a flow rate of 5 ml/min, where solvent A was 0.05% TFA/water and solvent B was 0.05% TFA/acetonitrile. The absorbance was recorded at 210 nm. The purity and authenticity of the peptides were examined by means of analytical HPLC, amino acid analysis, and mass spectrometry. Analytical HPLC was done on an Aquapore RP-300,  $\text{C}_8$  reversed-phase column ( $220 \times 4.6$  mm I.D., Brownlee Labs, Santa Clara, CA). Purified peptides were hydrolyzed with 6N HCl with 0.1% phenol in sealed, evacuated tubes at  $110^{\circ}\text{C}$  for 22 hours, and amino acid analyses were performed on a Beckman System 6300 High Performance Automatic Analyzer (Beckman, Palo Alto, CA). The molecular weight and purity of the peptide was confirmed with a Biolon 20 Plasma Desorption Time of Flight mass spectrometer (Uppsala, Sweden).

EXAMPLE 5COMPETITIVE ELISA SCREEN OF INDIVIDUAL PEPTIDES

Competitive ELISA was performed to determine the ability of the hexapeptides synthesized in inhibiting antibody binding to the peptide antigen. The assay was  
5 similar to the one detailed in Example 3, except that conventional 10 mM PBS was used instead of 50 mM PBS, and concentrations of the hexapeptides used during the incubation were varied.

This enabled the construction of a competitive binding curve from which  $I_{50}$  values could be determined. The  $I_{50}$  values were calculated using the software TableCurve  
10 (Jandel Scientific Inc., CA).

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- 5 (i) APPLICANT: S.P.I. Synthetic Peptides Incorporated
- (ii) TITLE OF INVENTION: Combinatorial Library Composition and Method
- 10 (iii) NUMBER OF SEQUENCES: 35
- (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: Dehlinger & Associates
- (B) STREET: 350 Cambridge Avenue, Suite 250
- 15 (C) CITY: Palo Alto
- (D) STATE: CA
- (E) COUNTRY: USA
- (F) ZIP: 94306
- 20 (v) COMPUTER READABLE FORM:
- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- 25 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- (B) FILING DATE: 13-JUN-1995
- (C) CLASSIFICATION:
- 30 (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/260,199
- (B) FILING DATE: 15-JUN-1994
- 35 (viii) ATTORNEY/AGENT INFORMATION:
- (A) NAME: Sholtz, Charles K.
- (B) REGISTRATION NUMBER: 38,615
- (C) REFERENCE/DOCKET NUMBER: 8900-0008.41
- 40 (ix) TELECOMMUNICATION INFORMATION:
- (A) TELEPHONE: (415) 324-0880
- (B) TELEFAX: (415) 324-0960

## (2) INFORMATION FOR SEQ ID NO:1:

27

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

5

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

10

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: Pseudomonas aeruginosa strain K  
(PAK) peptide

15

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

20

Lys Cys Thr Ser Asp Gln Asp Glu Gln Phe Ile Pro Lys Gly Cys Ser  
1                      5                      10                      15

Lys

25

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 7 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

30

## (ii) MOLECULE TYPE: peptide

35

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

45

Asp Glu Gln Phe Ile Pro Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:3:

5

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
10 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

15

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

20

(C) INDIVIDUAL ISOLATE: EQFIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

25 Glu Gln Phe Ile Pro Lys  
1 5

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
30 (A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

35

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

40

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: QQFIPK peptide

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

29

Gln Gln Phe Ile Pro Lys

1

5

## (2) INFORMATION FOR SEQ ID NO:5:

5

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

10

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

15

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: QAFIPK peptide

20

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Gln Ala Phe Ile Pro Lys

25

1

5

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

35

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

45

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

30

Ala Phe Phe Ile Pro Lys

1 5

## (2) INFORMATION FOR SEQ ID NO:7:

5

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

10

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

15

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: GQFIPK peptide

20

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gly Gln Phe Ile Pro Lys

25

1 5

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

35

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

40

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQFIIG peptide

45

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:



31

Glu Gln Phe Ile Ile Gly  
1 5

## (2) INFORMATION FOR SEQ ID NO:9:

5

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

10

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

15

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: QQFIIG peptide

20

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Gln Gln Phe Ile Ile Gly  
1 5

25

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

30

## (ii) MOLECULE TYPE: peptide

35

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## x1. SEQUENCE DESCRIPTION: SEQ ID NO:10:

32

Gln Ala Phe Ile Ile Gly  
1 5

## (2) INFORMATION FOR SEQ ID NO:11:

5

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

10

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

15

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: AFFIIG peptide

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Ala Phe Phe Ile Ile Gly  
1 5

25

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

30

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

35

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

40

(C) INDIVIDUAL ISOLATE: GQFIIG peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gly Gln Phe Ile Ile Gly  
1 5

45

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: peptide

10

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

15

## (vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQFIPL peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

20

Glu Gln Phe Ile Pro Leu  
1 5

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

30

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

35

## (vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: QQFIPL peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

## (2) INFORMATION FOR SEQ ID NO:15:

45

34

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 6 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: single  
5      (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO  
10

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
    (C) INDIVIDUAL ISOLATE: QAFIPL peptide  
15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Gln Ala Phe Ile Pro Leu  
1                      5  
20

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 6 amino acids  
    (B) TYPE: amino acid  
25      (C) STRANDEDNESS: single  
    (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO  
30

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
    (C) INDIVIDUAL ISOLATE: AFFIPL peptide  
35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Ala Phe Phe Ile Pro Leu  
1                      5  
40

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:  
45      (A) LENGTH: 6 amino acids

35

(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

5 (ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

10

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: GQFIPL peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

15

Gly Gln Phe Ile Pro Leu

1

5

(2) INFORMATION FOR SEQ ID NO:18:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

25

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

30

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: DQFIPK peptide

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Asp Gln Phe Ile Pro Lys

1

5

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

45

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

5 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

10 (C) INDIVIDUAL ISOLATE: ENFIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Glu Asn Phe Ile Pro Lys

15 1 5

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

25 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

30 (vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQYIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

35 Glu Gln Tyr Ile Pro Lys

1 5

(2) INFORMATION FOR SEQ ID NO:21:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

45

37

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

5 (iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQWIPK peptide

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Glu Gln Trp Ile Pro Lys

1 5

15 (2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

25

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQFMPK peptide

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Glu Gln Phe Met Pro Lys

1 5

35

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

40

(ii) MOLECULE TYPE: peptide

45

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQFVPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

10

Glu Gln Phe Val Pro Lys

1

5

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

20

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQFIPH peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

30

Glu Gln Phe Ile Pro His

1

5

(2) INFORMATION FOR SEQ ID NO:25:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

40

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

45



(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: EQFIPR peptide

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Glu Gln Phe Ile Pro Arg

1

5

10

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

15

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

20

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

25

(C) INDIVIDUAL ISOLATE: NQFIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Asn Gln Phe Ile Pro Lys

30

1

5

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

35

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

40

(iv) ANTI-SENSE: NO

45

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: QNFIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Gln Asn Phe Ile Pro Lys

1 5

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: NAFIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Asn Ala Phe Ile Pro Lys

1 5

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: GNFIPK peptide

41

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

Gly Asn Phe Ile Pro Lys  
1 5

5

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4 amino acids

(B) TYPE: amino acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

15 (iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

20 (C) INDIVIDUAL ISOLATE: FIPK peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

Phe Ile Pro Lys  
1

25

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

35 (ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

45

42

Gly Ala Tyr Ser His Gly  
1 5

## (2) INFORMATION FOR SEQ ID NO:32:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

10

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

15

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: GAYTHA peptide

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Gly Ala Tyr Thr His Ala  
1 5

25

## (2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
30 (C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: peptide

35

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

40

(C) INDIVIDUAL ISOLATE: VLYTKA peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

Val Leu Tyr Thr Lys Ala  
1 5

45

## (2) INFORMATION FOR SEQ ID NO:34:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: peptide

10

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

15

## (vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: AAWTKG peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

20

Ala Ala Trp Thr Lys Gly

1

5

## (2) INFORMATION FOR SEQ ID NO:35:

25

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: unknown

30

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

35

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: VLYSHG peptide

## IT IS CLAIMED:

1. A method of generating an oligomer compound capable of interacting specifically with a selected macromolecular ligand, comprising
  - 5 (a) reacting the ligand with each of a set of combinatorial oligomer libraries in which one or more selected subunit positions in the library oligomers have one of substantially all possible different subunits in each of the selected positions, and the remaining one or more subunit positions in each library include substantially all possible combinations of the different subunits,
  - 10 (b) selecting from the set of libraries, one or more libraries which show an elevated level of interaction with the ligand, thereby identifying one or more high-affinity subunit compositions corresponding to the selected subunit positions,
  - (c) repeating steps (a) and (b) with another group of selected one or more subunit positions, until substantially all of the subunit positions in the libraries oligomers have  
15 been considered,
  - (d) reacting the ligand with a permutation library of oligomers formed of placing the highest-affinity compositions identified in steps (a)-(c) in each of the associated oligomer positions, and
  - (e) selecting one or more oligomers from the permutation library which  
20 show the greatest level of interaction with said ligand.
2. The method of claim 1, for use in identifying a polypeptide oligomer capable of interacting specifically with a selected macromolecular ligand, wherein the subunits are representative amino acids that display the basic physico-chemical properties associated with  
25 naturally occurring amino acids, but exclude many of these naturally occurring amino acids.
3. The method of claim 2, wherein the subunits include at least one from each of the groups consisting of (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met, and Cys (h) Pro, (i) Gln and Asn, and (j) Ser and  
30 Thr.
4. The method of claim 1, for use in selecting a hexapeptide oligomer having three N-terminal and three C-terminal positions, wherein one set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three N-terminal positions,

and a second set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three C-terminal positions.

5        5.        The method of claim 4, wherein the subunits include at least one from each of the groups consisting of (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met, and Cys (h) Pro, (i) Gln and Asn, and (j) Ser and Thr, and each of said sets of libraries includes between about 500-2,000 libraries.

10       6.        The method of claim 1, for use in selecting a hexapeptide oligomer having two N-terminal, two intermediate, and two C-terminal positions, wherein one set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two N-terminal positions, a second set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two intermediate positions, and a third set of combinatorial libraries contains a library for each of the possible permutations of amino acids  
15       in the two C-terminal positions.

20       7.        The method of claim 6, wherein the subunits include at least one from each of the groups consisting of (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met, and Cys (h) Pro, (i) Gln and Asn, and (j) Ser and Thr.

25       8.        The method of claim 1, for use in identifying a oligomer having an upstream subunit position, and intermediate subunit position, and a downstream subunit position, where each subunit can be filled with substantially more than 20 different subunits, wherein one set of combinatorial libraries contains a library for each of the possible subunits in the upstream subunit position, a second set of combinatorial libraries contains a library for each of the possible subunits in the intermediate subunit position, and a third set of combinatorial libraries contains a library for each of the possible subunits in the downstream subunit position.

30       9.        The method of claim 8, wherein each subunit position can be filled by at least

10       10.       For use in selecting an oligomer compound capable of interacting specifically with a selected macromolecular ligand, a combinatorial library composition comprising

a first set of combinatorial oligomer libraries in which one or more selected subunit positions in the library oligomers have one of substantially all possible different subunits in each of the selected positions, and the remaining one or more subunit positions in each library include substantially all possible combinations of the different subunits, and

5 a second set of combinatorial oligomer libraries in which one or more different selected subunit positions in the library oligomers have one of substantially all possible different subunits in each of the selected positions, and the remaining one or more subunit positions in each library include substantially all possible combinations of the different subunits.

10 11. The composition of claim 10, for use in identifying a polypeptide oligomer capable of interacting specifically with a selected macromolecular ligand, wherein the subunits are representative amino acids that display the basic physico-chemical properties associated with naturally occurring amino acids, but exclude many of these naturally occurring amino acids.

15 12. The composition of claim 11, wherein the subunits include at least one from each of the groups consisting of (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met, and Cys (h) Pro, (i) Gln and Asn, and (j) Ser and Thr.

20 13. The composition of claim 10, for use in selecting a hexapeptide oligomer having three N-terminal and three C-terminal positions, wherein the first set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three N-terminal positions, and the second set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the three C-terminal positions.

25 14. The composition of claim 13, wherein the subunits include at least one from each of the groups consisting of (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met, and Cys (h) Pro, (i) Gln and Asn, and (j) Ser and Thr, and each of said sets of libraries includes between about 500-2,000 libraries.


30 15. The composition of claim 10, for use in selecting a hexapeptide oligomer having two N-terminal, two intermediate, and two C-terminal positions, wherein the first set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two N-terminal positions, and the second set of combinatorial libraries contains a



library for each of the possible permutations of amino acids in the two intermediate positions, and which further includes a third set of combinatorial libraries contains a library for each of the possible permutations of amino acids in the two C-terminal positions.

- 5           16.     The composition of claim 15, wherein the subunits include at least one from each of the groups consisting of (a) Ala, (b) Glu and Asp, (c) Phe, Tyr, and Trp, (d) Gly, (e) Ile and Val, (f) Lys, His, and Arg, (g) Leu, Met, and Cys (h) Pro, (i) Gln and Asn, and (j) Ser and Thr.
- 10           17.     The composition of claim 10, for use in identifying a oligomer having an upstream subunit position, and intermediate subunit position, and a downstream subunit position, where each subunit can be filled with substantially more than 20 different subunits, wherein the first set of combinatorial libraries contains a library for each of the possible subunits in the upstream subunit position and the second set of combinatorial libraries contains  
15 a library for each of the possible subunits in the intermediate subunit position, and which further includes a third set of combinatorial libraries contains a library for each of the possible subunits in the downstream subunit position.
18.     The method of claim 17, wherein each subunit position can be filled by at least  
20 about 50 different subunits.

1/10



$AAA \ X_4 \ X_5 \ X_6$   
 $AAR \ X_4 \ X_5 \ X_6$   
 $ARA \ X_4 \ X_5 \ X_6$

$VVV \ X_4 \ X_5 \ X_6$


**Fig. 2A**

$X_1 \ X_2 \ X_3 \ AAA$   
 $X_1 \ X_2 \ X_3 \ AAR$   
 $X_1 \ X_2 \ X_3 \ ARA$

$X_1 \ X_2 \ X_3 \ VVV$

**Fig. 2B**

2/10



AA X<sub>3</sub> X<sub>4</sub> X<sub>5</sub> X<sub>6</sub>  
 AR X<sub>3</sub> X<sub>4</sub> X<sub>5</sub> X<sub>6</sub>  
 RA X<sub>3</sub> X<sub>4</sub> X<sub>5</sub> X<sub>6</sub>

VV X<sub>3</sub> X<sub>4</sub> X<sub>5</sub> X<sub>6</sub>

**Fig. 4A**

X<sub>1</sub> X<sub>2</sub> AA X<sub>5</sub> X<sub>6</sub>  
 X<sub>1</sub> X<sub>2</sub> AR X<sub>5</sub> X<sub>6</sub>  
 X<sub>1</sub> X<sub>2</sub> RA X<sub>5</sub> X<sub>6</sub>

X<sub>1</sub> X<sub>2</sub> X<sub>3</sub> X<sub>4</sub> AA  
 X<sub>1</sub> X<sub>2</sub> X<sub>3</sub> X<sub>4</sub> AR  
 X<sub>1</sub> X<sub>2</sub> X<sub>3</sub> X<sub>4</sub> RA

X<sub>1</sub> X<sub>2</sub> VV X<sub>5</sub> X<sub>6</sub>

**Fig. 4B**

X<sub>1</sub> X<sub>2</sub> X<sub>3</sub> X<sub>4</sub> VV

**Fig. 4C**

3/10

$s_1$   $s_2$   $s_3$   
| | |  
 $R_i$   $R_j$   $R_k$

**Fig. 5A**




$s_1(R_1)$	$s_2(R_j)$	$s_3(R_k)$	$s_1(R_i)$	$s_2(R_j)$	$s_3(R_1)$
$s_1(R_2)$	$s_2(R_j)$	$s_3(R_k)$	$s_1(R_i)$	$s_2(R_2)$	$s_3(R_2)$
.	.	.	.	.	.
$s_1(R_n)$	$s_2(R_j)$	$s_3(R_k)$	$s_1(R_i)$	$s_2(R_n)$	$s_3(R_n)$

**Fig. 5B**


**Fig. 5C**

**Fig. 5D**


4/10

**Fig. 6A****Fig. 6B****Fig. 7**

5/10



X1 X2 O3 O4 X5 X6 → FI

**Fig. 8B****Fig. 8C**

SEQ ID:3: EQFIPK\*  
 SEQ ID:4: QQFIPK\*  
 SEQ ID:5: QAFIPK\*  
 SEQ ID:6: AFFIPK  
 SEQ ID:7: GQFIPK  
 SEQ ID:8: EQFIIG

**Fig. 9****Fig. 10**

SEQ ID:18: DQFIPK\*  
 SEQ ID:19: ENFIPK  
 SEQ ID:20: EQYIPK  
 SEQ ID:21: EQWIPK\*  
 SEQ ID:22: EQFMPK\*  
 SEQ ID:23: EQFVPK\*  
 SEQ ID:24: EQFIPH  
 SEQ ID:25: EQFIPR  
 SEQ ID:26: NQFIPK  
 SEQ ID:27: QNFIPK  
 SEQ ID:28: NAFIPK  
 SEQ ID:29: GNFIPK

6/10




Fig. 11A

7/10





Fig. 11B






8/10




**Fig. 11C**



9/10

**Fig. 12****Fig. 13A****Fig. 13B**

10/10



58

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/IB 95/00560

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07K1/04 G01N33/53

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category \* Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No.

X	WO,A,86 00991 (COMMONWEALTH SERUM LABORATORIES COMMISSION) 13 February 1986 * page 10; claims 1-5 *	1-18
---	--	------

P,X	METHODS, vol. 6, 1994 pages 404-410, WONG, W.Y. ET AL 'Representative combinatorial peptide libraries: an approach to reduce both synthesis and screening efforts' * whole disclosure *	1-18
-----	--	------

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

22 September 1995

3 1. 10. 95

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Hermann, R

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 95/00560

(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NATURE, vol. 354, 1991 pages 84-86, HOUGHTEN, R.A. ET AL. 'Generation and use of synthetic peptide combinatorial libraries for basic research and drug discovery' cited in the application * whole disclosure *</p> <p>---</p>	1-18
A	<p>INT. J. PEPTIDE PROTEIN RES., vol. 37, 1991 pages 487-493, FURKA, A. 'General methods for rapid synthesis of multicomponent peptide mixtures' cited in the application * whole disclosure *</p> <p>---</p>	1-18
X	<p>WO,A,92 09300 (ITEREX PHARMACEUTICALS LTD) 11 June 1992 * claims; page 55, line 24 to page 56, line 19 *</p> <p>-----</p>	1-18

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 95/00560

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-8600991	13-02-86	AU-B- 592311	11-01-90
		AU-B- 4533985	30-01-86
		CA-A- 1255586	13-06-89
		EP-A- 0190205	13-08-86
		JP-T- 61502839	04-12-86
		US-A- 5194392	16-03-93
-----			
WO-A-9209300	11-06-92	CA-A- 2090860	22-05-92
		EP-A- 0558671	08-09-93
		JP-T- 6507378	25-08-94
-----			